

Review Article

A place for palm fruit oil to eliminate vitamin A deficiency*

A J Spinnler Benadé DSc

Nutritional Intervention Research Unit, Medical Research Council, Parow, South Africa

There is general consensus that food-based approaches are viable and sustainable options for addressing vitamin A deficiency in populations. One such example is the fortification of food which, if properly monitored, could make a significant contribution towards improving the vitamin A status of populations throughout the world. Red palm fruit oil (RPO) with its high content of natural carotenoids, lends itself exceptionally well to this purpose at both household and commercial level. Results are now available from several feeding trials incorporating RPO into diets at household level or into commercially manufactured products. RPO in the maternal diet was shown to improve the vitamin A status of lactating mothers and their infants. Consumption of RPO incorporated in a sweet snack or biscuits significantly improved plasma retinol concentrations in children with subclinical vitamin A deficiency. There is evidence that if only 35-50% of the recommended daily intake for vitamin A were to be provided by RPO, it may be sufficient to prevent vitamin A deficiency (hypovitaminosis A). Red palm oil has a highly bioconvertible form of alpha- and beta-carotene, a long shelf life, and a higher cost/benefit ratio when compared to other approaches such as high-dose-vitamin A supplements and fortification of foods with retinyl ester fortificants. Consumption of RPO is safe and cannot produce hypervitaminosis A. Considering all the current information about RPO, the initiation of food-based interventions involving its use in developing countries with an endemic vitamin A deficiency problem, appears to be a logical choice.

Key Words: red palm oil, vitamin A deficiency, fortification, developing countries, food-based approaches

Introduction

According to a recent report, the global prevalence for clinical vitamin A deficiency for preschool children in 1995 was estimated to be 1.2%.¹ For subclinical vitamin A deficiency, on a global basis, it is estimated that 140 million preschoolers are affected by subclinical vitamin A deficiency.¹ The highest prevalences of both clinical and subclinical vitamin A deficiency occur in south Asia and Sub-Saharan Africa where 30-40% of preschool children are at an increased risk of ill health and death because of this deficiency. Of the 140 million preschoolers world-wide who are affected by subclinical vitamin A deficiency, 60 million live in south Asia and 40 million in Sub-Saharan Africa.¹

Several strategies for alleviation of vitamin A deficiency have been proposed. These include: 1) high-dose vitamin A capsule supplementation, 2) food fortification, and 3) food diversification and nutritional education. Although supplementation with high-dose vitamin A capsules, when targeted exclusively at the young child population who is at a higher risk for hypovitaminosis A, is the most cost-effective intervention, there is global consensus that food-based strategies are viable and sustainable options for addressing micronutrient deficiencies in populations. Fortification of food, when universal coverage is the goal, appears to be the least expensive strategy.

Addressing vitamin A deficiency through food fortification

Retinyl palmitate is the vitamin A fortificant most commonly used in food fortification and has been used to fortify sugar in

Guatemala², monosodium glutamate in the Philippines and Indonesia^{3,4}, and wheat flour in the Philippines.⁵ The Guate-malan sugar fortification programme was evaluated over a period of two years, and documented a significant improvement in vitamin A status in preschool children, especially in those with low serum retinol levels at baseline.² In the intestinal lumen retinyl palmitate is hydrolysed to retinol and absorbed by carrier-mediated diffusion or simple diffusion. The absorption of vitamin A is considered to be a non-saturable mechanism,⁶ and uncontrolled and excessive intake may therefore lead to vitamin A toxicity.

Beta-carotene, a precursor of vitamin A, has been used as a safe colourant in foods and beverages for many years; it turns out that it also has the potential for use as a vitamin A fortificant. The amount of beta-carotene that is converted to vitamin A in the intestinal mucosa depends on the vitamin A status of the individual⁷ which, when adequate, has an inhibitory effect on the enzyme that cleaves beta-carotene into retinal.⁸ The risk of vitamin A toxicity through overconsumption is therefore eliminated if beta-carotene is used as a vitamin A fortificant.

Correspondence address: Dr AJS Benadé, NIRU, MRC
PO Box 19070, Tygerberg 7505, South Africa.
Tel: +27 21 9380283; Fax: +27 21 9380321
Email: spinnler.benade@mrc.ac.za

Accepted 30 June 2003

*Presented at Symposium on "North & West African Foods and Health" February 8th 2003, Marrakech, Morocco

Synthetic beta-carotene was used as a fortificant in a wafer given to lactating women in Indonesia, and proved to be more effective in improving the vitamin A status of these women than the same amount of beta-carotene provided by dark green leafy vegetables.⁹ Synthetic beta-carotene has also been used to fortify soybean cooking oil in Brazil, and studies have shown that the bioavailability of beta-carotene in the oil was not influenced by heat treatment.¹⁰

Characteristics of crude and refined red palm fruit oil

Oil from the African palm (genus *Elaeis*) is a rich natural source of provitamin A carotenoids and can be used as an alternative vitamin A fortificant. Crude palm oil is obtained from the mesocarp of the African palm fruit¹¹ and contains 500-700 ppm of carotenoids of which 56% and 35% are comprised of beta- and alpha-carotene, respectively.¹² Although the proportion of these two carotenes is similar in palm oil and carrots, red palm fruit oil (RPO) contains about 15 times more carotenes than that present in the same weight of carrots, and 44 times that of leafy vegetables.¹³

Crude palm oil has a very strong taste and odour and is not very stable, due to free fatty acids and other impurities. It can be refined, however, to a palatable red oil that is odourless, tasteless and very stable, but in which 80% of the carotenes and vitamin E originally present in the crude palm oil are retained.¹¹ The process developed in Malaysia to produce deodorised and deacidified red palm oil involves two stages, namely pretreatment of the crude palm oil followed by decolourisation and deodorisation by molecular distillation.¹¹ The product has been given the name Carotino.

Value of red palm fruit oil for combatting vitamin A deficiency

Results from several efficacy trials have confirmed the value of red palm fruit oil in improving vitamin A status. A few of them from various nations and foci of interest are reviewed below.

Red palm oil in the maternal diet - vitamin A status of mothers and their nursing infants

Canfield *et al.*,¹⁴ studied the effect of refined RPO supplementation in a population of lactating mothers and their nursing infants in Honduras. Mothers received either a placebo (N=8), 15 mg beta-carotene (N=20), or 15 mg beta-carotene (N=26) as RPO, on six occasions over a period of two weeks. At the beginning of the study, approximately 60% of the infants had serum vitamin A concentrations indicative of borderline vitamin A deficiency (<0.7 µmol). At the end of the supplementation period, serum retinol of infants of mothers receiving beta-carotene or palm oil was significantly increased relative to the placebo group. The authors conclude that RPO in the diet of lactating mothers has the potential to enhance vitamin A status of their nursing infants and thereby enhance their resistance to morbidity due to respiratory and intestinal infections.

Effect of red palm oil on vitamin A status of school children

Manorama and co-workers¹⁵ reported results from three feeding trials on schoolchildren receiving sweet snacks containing red palm oil supplying the Indian recommended dietary allowances (2400µg) of beta-carotene. In the first trial, children of 7-9 years of age²⁴ received the snack for

two months, after which time significant increases were seen in serum retinol levels. Serum retinol levels increased from basal 0.86 ± 0.13 to 1.89 ± 0.023 µmol/L. These changes were similar to those seen in children receiving 600µg of vitamin A daily. Liver reserves of retinol, as determined by the Modified Relative Dose Response Test, indicated a saturation of liver reserves of retinol.

In the second trial, one group of schoolchildren (N=18) again received the red palm oil snack for one month, while a second group (N=18) received 100 000 IU of synthetic vitamin A as a single massive dose. Serum retinol concentrations increased significantly in both groups, 1.40 ± 0.05 to 1.76 ± 0.09 and 0.95 ± 0.05 to 1.85 ± 0.008 µmol/L in the vitamin A and RPO groups, respectively. In the third trial, three groups of children (12 per group) received either a single massive vitamin A dose of 100 000 IU as vitamin A palmitate, 4 g red palm oil for 30 days providing 50 000 IU vitamin A, or 8 g red palm oil for 30 days providing 100 000 IU vitamin A.¹⁵

All treatment groups showed significant increases in serum retinol levels measured 30 days after the single vitamin A dose or daily supplementation with palm oil. Although all children had serum retinol levels < 0.7 µmol/L at baseline, none had levels < 0.7 µmol/L after 30 days. Six months after cessation of supplementation with RPO, mean serum retinol levels were maintained at > 0.7 µmol/L in the vitamin A and 8 g per day RPO groups. Mean values in the 4 g red palm oil group were lower than 0.7 µmol/L six months after cessation of supplementation. Even when children were fed 50% of the Indian recommendations from the RPO snack, normal serum levels of retinol were maintained six months after cessation of supplementation. However, 33% of children in this group had retinol levels < 0.7 µmol/L at the end of the 6-month period.

Red palm oil-based baking fat as a food fortificant for combatting vitamin A deficiency

Van Stuijvenberg *et al.*,¹⁶ evaluated the response of serum retinol concentrations in rural schoolchildren (N=115) 6-11 consuming biscuits fortified with beta-carotene at 50% of their RDA for retinol. From these feeding trials, it was concluded that beta-carotene can be used as a food fortificant for combatting vitamin A deficiency.

The use of beta-carotene in the baking process was fraught with technical difficulties such as weighing errors in the preparation of the premix, instability of beta-carotene when exposed to light and air, and a wide variation in the beta-carotene content of the biscuits baked at different sites. Furthermore the baking fat used in the baking process was hydrogenated fat containing large amounts of *trans* fatty acids which are known to be associated with negative health effects.¹⁷

The manufacturers of Carotino were subsequently requested to develop a baking fat containing all the nutritional properties of Carotino oil, and without *trans* fatty acids. This led to the development of Carotino baking fat which was subsequently evaluated in terms of its value as a food fortificant for combatting vitamin A deficiency. In a placebo controlled trial,¹⁸ schoolchildren 5-11 of age were randomly assigned to three different treatment categories, namely: 1) a biscuit containing no added beta-carotene (N=137); 2) a biscuit with synthetic beta-carotene (N=130);

3) a biscuit with the refined red palm oil shortening as a source of beta-carotene (N=133). Biscuits were distributed daily during the school week for three months. The synthetic beta-carotene and RPO biscuits were designed to provide a similar amount of beta-carotene, providing approximately 34% of the RDA for 7-10-year-old children, assuming a conversion factor of beta-carotene to retinol of 6:1.¹⁹ There was a significant improvement compared to the control group in both the synthetic beta-carotene and the RPO group. The effects of the two treatments were found to be equivalent. The percentage of children with serum retinol concentrations below 15 µg/dL dropped from 17.5% to 13.1% in the control group, from 17.7% to 4.6% in the synthetic β-carotene group, and from 15.8% to 6.8% in the RPO group (the authors used <15 µg/dL as a cut-off value, instead of the normal cut-off of <20 µg/dL, because the majority of the values for this population lay around 20 µg/dL and, as such, this cut-off value was not sensitive enough for illustrating differences in prevalence changes between the control and intervention groups) (Table 1).

Potential of Carotino shortening as food fortificant for vitamin A in the baking industry

It is now well established that Carotino baking fat lends

Table 1. Serum retinol (mg/dL) in primary schoolchildren before and after three months of intervention with biscuits fortified with synthetic β-carotene or red palm oil biscuit

Measurement	Control biscuit (N = 137)	β-Carotene-fortified biscuit (N = 130)	Red palm oil biscuit (N = 133)
Baseline	20.6 (5.8) ^a 17.5% ^b	20.4 (6.0) 17.7%	20.8 (7.0) 15.8%
3 months	21.6 (5.9) 13.1%	24.4 (5.6) 4.6% ^c	24.0 (6.7) 6.8% ^c

^aMean (SD); ^b[percentage of children with low serum retinol levels] using a cut-off of <15 mg/dL; ^cp<.005 compared with change in control group (ANOVA); Reproduced from 18.

itself exceptionally well to application as a food fortificant in the food industry. Calculations based on the carotene content of Carotino (assuming that 6 µg of dietary all-*trans*-beta-carotene equals 1 µg of all-*trans*-retinol) and the fat content of a variety of bakery products showed that Carotino shortening could supply between 46% and 70% of the RDA for vitamin A²⁰ for children 7-10 of age (Table 2).

Additional benefits of Carotino shortening.

Apart from its potential as a food fortificant for vitamin A in the food and baking industry, incorporation of Carotino in a variety of consumer products offers several additional advantages.

Carotino contains no *trans* fatty acids

The health implications of *trans* fatty acid consumption are well established. However, hydrogenated fat, containing variable amounts of *trans* fatty acids, is extensively used in the baking industry. Because of the process used in the manufacture of Carotino baking fat, Carotino does not require hydrogenation, and is therefore free of *trans* fatty acids. *Trans* fatty acids are known to behave metabolically like saturated fat and are atherogenic.^{21,22}

Vitamin E activity of Carotino baking fat

Carotino baking fat is a rich source of tocopherols and tocotrienols which have different vitamin E activity and which are also natural antioxidants. They act as scavengers of the damaging oxygen-free radicals that are suggested to play a role in cellular aging, atherosclerosis and cancer.²³⁻²⁵

Carotenoid content of Carotino baking fat.

Carotino baking fat contains a broad spectrum of carotenoids,²⁶ some of which could serve as precursors of vitamin A while others have antioxidant properties.

Stability and safety of Carotino baking fat

Because of the high natural antioxidant content of Carotino baking fat, it does not require the addition of synthetic or other antioxidant to consumer products containing Carotino baking fat. Biscuits baked with Carotino fat proved to have a shelf life exceeding 6 months when stored at temperatures

Table 2. Retinol equivalents (RE) and tocopherol equivalents (TE) per portion of various food products

Product	Fat content (%)	Portion (g)	% RDA RE		% RDA TE	
			7-10 yr	>10 yr	7-10 yr	>10 yr
Carotino biscuits	16.0	45	62	54	37	26
Banana loaf	10.8	50	46	41	27	19
Bread, maize meal	11.3	50	48	42	29	20
Cake, home-made	11.8	50	48	42	29	20
Chocolate cake	12.9	50	55	48	33	23
Cookies, plain	13.4	20	23	20	14	10
Oat crunchies	23.4	25	50	44	29	20
Carrot cake	19.7	50	85	74	51	36
Fruit cake	12.9	50	55	48	33	23
Cookies, shortbread	27.2	20	46	41	27	19
Crackers	24.6	33	70	62	41	29
Gingerbread	12.6	50	54	47	33	23
Scone, whole wheat	12.6	50	54	47	33	23
Muffin, oat bran	9.7	50	41	37	24	17

Reproduced from 20, with addition

of 20-30°C in the dark. Close monitoring of school learners consuming Carotino biscuits over a period of one year revealed no negative side-effects on health (Benadé AJS: unpublished observation).

Quality control of beta-carotene content of biscuits baked with Carotino fat

Because the carotene content of Carotino fat is controlled at the point of manufacture, it is possible for different manufacturers to produce products with exceptional uniformity in terms of its beta-carotene content.

Conversion of beta-carotene from Carotino baking fat to retinol

It has been suggested that the conversion factor of beta-carotene in oil to retinol is 2:1.¹⁹ Based on this assumption, the amount of fat in products could be adjusted accordingly, which will not only reduce cost, but will also reduce the intensity of the yellow colour in products prepared with Carotino.

Summary and Conclusions

Results from several studies have confirmed the value of red palm oil as a food fortificant for combatting vitamin A deficiency. Based on current knowledge regarding the composition of red palm oil and the potential health benefits of these components, the value of RPO as a functional food justifies further investigation. Although the colour of the oil may hamper its acceptability by consumers as a day-to-day cooking oil, new innovative products containing RPO could overcome this problem.

Acknowledgement

I should like to thank the Malaysian Palm Oil Promotion Council for their financial support to attend the Symposium on "North and West African Foods and Health".

References

1. The Micronutrient Initiative. The micronutrient report: current progress and trends in the control of vitamin A, iodine, and iron deficiencies. Ottawa, Canada: The Micronutrient Initiative, International Development Research Center, Ottawa, Canada, 2001.
2. Arroyave G, Mejia LA, Aguilar JP. The effect of vitamin A fortification of sugar on the serum vitamin A levels of preschool Guatemalan children: a longitudinal evaluation. *Am J Clin Nutr* 1981; 34: 41-49.
3. Solon FS, Latham MC, Guirriec R, Florentino R, Williamson DF, Aguilar J. Fortification of MSG with vitamin A: the Philippines experience. *Food Technology* 1985; 39: 71-79.
4. Muhilal, Murdiana A, Azio I, Saidin S, Jahari AB, Karyadi D. Vitamin A-fortified monosodium glutamate and vitamin A status: a controlled field trial. *Am J Clin Nutr* 1988; 48: 1265-1270.
5. Solon FS, Klemm RDW, Sanchez L, Darnton-Hill I, Craft NE, Christian P, West KP Jr. Efficacy of a vitamin A-fortified wheat-flour bun on the vitamin A status of Filipino schoolchildren. *Am J Clin Nutr* 2000; 72: 738-744.
6. Biesalski HK. Bioavailability of vitamin A. *Eur J Clin Nutr* 1997; 51: 571-575.
7. Ribaya-Mercado JD, Solon FS, Solon MA, Cabal-Barza MA, Perfecto CS, Tang G, Solon JAA, Fjeld CR, Russel RM. Bioconversion of plant carotenoids to vitamin A in Filipino school-aged children varies inversely with vitamin A status. *Am J Clin Nutr* 2000; 72: 455-465.
8. Villard L, Bates CJ. Carotene dioxygenase activity in rat intestine: effects of vitamin A deficiency and of pregnancy. *Br J Nutr* 1986; 56: 115-122.
9. de Pee S, West CE, Muhilal, Karyadi D, Hautvast JGAJ. Lack of improvement in vitamin A status with increased consumption of dark-green leafy vegetables. *Lancet* 1995; 346: 75-81.
10. Dutra-de-Oliveira JE, Fávoro RMD, Junqueira-Franco MVM, Carvalho CG, Jordão AA, Vannucchi H. Effect of heat treatment on the biological value of β -carotene added to soybean cooking oil in rats. *Int J Food Sci Nutr* 1998; 49: 205-210.
11. Nagendran B, Unnithan UR, Choo YM, Sundram K. Characteristics of red palm oil, a carotene- and vitamin E-rich refined oil for food uses. *Food Nutr Bull* 2000; 21: 189-194.
12. An Endorsement on Health, Nutrition and Palm Oil. Selangor, Malaysia: Malaysian Palm Oil Promotion Council, 2000.
13. Scrimshaw NS. Nutritional potential of red palm oil for combating vitamin A deficiency. *Food Nutr Bull* 2000; 21: 195-201.
14. Canfield LM, Liu Y, de Kaminsky R, Castillo C, Zavala G, Garner C, Pagoaga E. Supplementation of mothers with red palm oil increases infant vitamin A status. Proceedings of the PORIM International Palm Oil Congress (PIPOC), Kuala Lumpur, Malaysia, 1996.
15. Manorama R, Sarita M, Rukmini C. Red palm oil for combating vitamin A deficiency. *Asia Pac J Clin Nutr* 1997; 6: 56-59.
16. Van Stuijvenberg ME, Kvalsvig JD, Faber M, Kruger M, Kenoyer DG, Benadé AJS. Effect of iron-, iodine- and β -carotene-fortified biscuits on the micronutrient status of primary school children: a randomized controlled trial. *Am J Clin Nutr* 1999; 69: 497-503.
17. Willett WC, Ascherio A. Trans-fatty acids: are the effects only marginal? *Am J Pub Health* 1994; 84: 722-724.
18. Van Stuijvenberg ME, Benadé AJS. South African experience with the use of red palm oil to improve the vitamin A status of primary schoolchildren. *Food Nutr Bull* 2000; 21: 212-214.
19. Trumbo P, Yates AA, Schlicker S, Pood M. Dietary reference intakes: vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. *J Am Diet Assoc* 2001; 101: 294-301.
20. Benadé AJS. The potential of red palm oil-based shortening as a food fortificant for vitamin A in the baking industry. *Food Nutr Bull* 2001; 22: 416-418.
21. Mensink RPM, Katan MB. Effect of dietary *trans* fatty acids on high-density and low-density lipoprotein cholesterol levels in healthy subjects. *N Engl J Med* 1990; 323: 439-445.
22. Judd JT, Clevidence BA, Muesing BA, Wittes J, Sunkin ME, Podczasy JJ. Dietary *trans* fatty acids: effects on plasma lipids and lipoproteins of healthy men and women. *Am J Clin Nutr* 1994; 59: 861-868.
23. Pryor WA, Ames BN, Saul RC, McCord JM, Harman D. Oxygen radicals and human disease. *Ann Intern Med* 1987; 107: 526-545.
24. Jozwiak Z, Jasnowska B. Changes in oxygen metabolizing enzymes and lipid peroxidation in human erythrocytes as a function of age of donor. *Mech Ageing Dev* 1985; 32: 77-83.
25. Van Pappel G, Kardinal A, Princen H, Kok FJ. Antioxidants and coronary heart disease. *Ann Med* 1994; 26: 429-434.
26. Gellenbeck KW. Carotenoids: more than just β -carotene. *Asian J Clin Nutr* 1998; 7: 272-281.